

4150 Woodlands Pkwy, Suite B  
Palm Harbor, FL 34685  
(727) 772-1452



2137 Little Rd  
Trinity, FL 34655  
(727) 372-6760

---

## SPITTING UP IN INFANTS (GE Reflux)

### What is spitting up?

Spitting up (also called reflux) is the effortless spitting up of one or two mouthfuls of stomach contents. Formula or breast milk just rolls out of the mouth, often with a burp. It usually happens during or shortly after feedings. It begins in the first weeks of life. Spitting up is harmless as long as your infant doesn't spit up large amounts that interfere with normal weight gain.

### What is the cause?

Spitting up results from poor closure of the valve (ring of muscle) at the upper end of the stomach. Spitting up is normal and harmless for over half of all babies. It becomes a problem if it causes poor weight gain (from spitting up large amounts), choking, or acid damage to the lower esophagus (esophagitis).

### How long does it last?

Spitting up improves with age. By 7 months of age, most reflux has decreased or is gone. The reasons for this are probably because the baby is old enough to sit up or is eating solid foods. By the time your baby has been walking for 3 months, even severe reflux should be totally cleared up.

### How can I take care of my child?

- **Feed smaller amounts.** Overfeeding always makes spitting up worse. If the stomach is filled to capacity, spitting up is more likely. Give your baby smaller amounts. Your baby doesn't have to finish a bottle. Wait at least 2 ½ hours between feedings because it takes that long for the stomach to empty itself.
- **Avoid pressure on your child's abdomen.** Avoid tight diapers; they put added pressure on the stomach. Don't put pressure on the stomach or play vigorously with him/her right after meals.
- **Burp your child to reduce spitting up.** Burp your baby two or three times during each feeding. Do it when he/she pauses and looks around; don't interrupt feeding rhythm in order to burp him/her. Keep in mind that burping is less important than giving smaller feedings and avoiding tight diapers.
- **Keep your child in a vertical position after meals.** After meals, try to keep your baby in an upright position using a frontpack, backpack, or swing for 30 minutes. When your infant is in an infant seat, keep him from getting scrunched up by putting a pad under his buttocks so he's more stretched out. After your child is over 6 months old, a jumpy seat or infant play station can be helpful for maintaining an upright posture after meals.
- **Use a proper sleep position.** Most infants with spitting up problems can sleep on their backs, the position recommended by the American Academy of Pediatrics to reduce the risk of SIDS. If the esophagus becomes irritated (esophagitis), we may recommend having your baby sleep on his/her right side. Try to elevate the head of the bed a bit. If your child is having breathing problems (choking or sleep apnea), please call the office.
- **Add rice cereal to formula.** If your infant still spits up large amounts after all the previous treatments have been tried, you can try thickening the formula with rice cereal. Add 1 level teaspoon of rice cereal to each ounce of formula. The nipple opening will need to be enlarged. **\*\*Please do not add rice cereal to your infant's formula without consulting us first.\*\***
- **Acid blockers or liquid antacids.** Children with severe reflux or symptoms of heartburn may need temporary reduction of stomach acid.

Your child's medication is \_\_\_\_\_ Directions:  
\_\_\_\_\_

### When should I call the office?

Call IMMEDIATELY if:

- You see blood in the spit-up material.
- The spitting up causes your child to choke or cough.

Call during office hours if:

- Your baby doesn't seem to improve with this approach.
- Your baby does not gain weight normally.
- You have other concerns or questions.